

Gasoline emissions dominate over diesel in formation of secondary organic aerosol mass

R. Bahreini,^{1,2} A. M. Middlebrook,² J. A. de Gouw,^{1,2} C. Warneke,^{1,2} M. Trainer,² C. A. Brock,² H. Stark,^{1,2,3} S. S. Brown,² W. P. Dube,^{1,2} J. B. Gilman,^{1,2} K. Hall,^{1,2,4} J. S. Holloway,^{1,2} W. C. Kuster,² A. E. Perring,^{1,2} A. S. H. Prevot,⁵ J. P. Schwarz,^{1,2} J. R. Spackman,^{1,2,6} S. Szidat,⁷ N. L. Wagner,^{1,2} R. J. Weber,⁸ P. Zotter,⁵ and D. D. Parrish²

Received 20 December 2011; revised 16 February 2012; accepted 26 February 2012; published 20 March 2012.

[1] Although laboratory experiments have shown that organic compounds in both gasoline fuel and diesel engine exhaust can form secondary organic aerosol (SOA), the fractional contribution from gasoline and diesel exhaust emissions to ambient SOA in urban environments is poorly known. Here we use airborne and ground-based measurements of organic aerosol (OA) in the Los Angeles (LA) Basin, California made during May and June 2010 to assess the amount of SOA formed from diesel emissions. Diesel emissions in the LA Basin vary between weekdays and weekends, with 54% lower diesel emissions on weekends. Despite this difference in source contributions, in air masses with similar degrees of photochemical processing, formation of OA is the same on weekends and weekdays, within the measurement uncertainties. This result indicates that the contribution from diesel emissions to SOA formation is zero within our uncertainties. Therefore, substantial reductions of SOA mass on local to global scales will be achieved by reducing gasoline vehicle emissions. **Citation:** Bahreini, R., et al. (2012), Gasoline emissions dominate over diesel in formation of secondary organic aerosol mass, *Geophys. Res. Lett.*, 39, L06805, doi:10.1029/2011GL050718.

1. Introduction

[2] Anthropogenic aerosol particles have adverse effects on air quality, visibility, and health as well as direct and indirect effects on climate [Monks *et al.*, 2009]. Among the common aerosol types, organic aerosol (OA) often contributes a large fraction of the submicron aerosol mass [Zhang *et al.*, 2007]. Laboratory experiments have indicated that OA can be formed from photooxidation of precursors present in gasoline fuel and diesel exhaust [Miracolo *et al.*, 2010; Odum *et al.*, 1997] and

from biogenic emissions [Griffin *et al.*, 1999]. However, the relative importance of these sources of OA in many regions is poorly understood [Fuzzi *et al.*, 2006; Monks *et al.*, 2009].

[3] Emissions from gasoline and diesel vehicles differ in their chemical characteristics [Kleeman *et al.*, 2008; Presto *et al.*, 2009] and their weekly pattern [Harley *et al.*, 2005; Marr and Harley, 2002a]. In the United States, emissions from gasoline vehicles dominate vehicular hydrocarbon and carbon monoxide (CO) emissions [Harley *et al.*, 2001; Marr and Harley, 2002b]. Emission factors of nitrogen oxides (NO_x), black carbon (BC), PM_{2.5} (particulate matter, less than 2.5 micron), and primary OA (POA) from diesel engines are, respectively, a factor of 13, 37, 20, and 13 higher than from gasoline, per mass of fuel used [Ban-Weiss *et al.*, 2008]. Gasoline and diesel fuel sale data show that diesel fuel consumption was only 17% of the total fuel consumption on weekdays during May–June 2010 in California (<http://www.boe.ca.gov/sptaxprog/spftrpts.htm>, see auxiliary material).¹ Because of the health hazards associated with primary particles emitted from diesel vehicles, regulations in California since 1998 are directed to control PM emissions from diesel vehicles (<http://www.arb.ca.gov/msprog/onroadhd/onroadhd.htm>, and links therein). Traffic counts have indicated that vehicular diesel emissions in California are lower on weekends compared to weekdays by 60–80% while only a 5–10% decrease in gasoline-powered vehicles is observed on the weekends [Harley *et al.*, 2005]. Effects of lower diesel emissions during weekends on NO_x, and subsequently on ozone, have been studied in many urban areas [Harley *et al.*, 2005; Marr and Harley, 2002a, 2002b; Murphy *et al.*, 2007; Pollack *et al.*, 2012]. These studies indicate that there is a large decrease in emissions of NO_x and only a small decrease in emissions of VOCs during the weekends. The combination of a more rapid ozone production at higher VOC/NO_x ratios in VOC-limited areas [Marr and Harley, 2002a; Pollack *et al.*, 2012] and less O₃ destruction by NO_x titration [Murphy *et al.*, 2007; Pollack *et al.*, 2012] leads to higher average ozone observed in urban areas over the weekend. Analysis of organic and elemental carbon (OC, EC, respectively) measurements in Southern California during summer and fall of 1995 [Harley *et al.*, 2005] and the IMPROVE (Interagency Monitoring of Protected Visual Environments) aerosol samples throughout the U.S. [Murphy *et al.*, 2008] indicate that fine particle EC has a pronounced weekly cycle, with minima occurring on Sunday, while OC concentrations are more constant.

¹Cooperative Institute for Research in Environmental Sciences, University of Colorado at Boulder, Boulder, Colorado, USA.

²Earth System Research Laboratory, National Oceanic and Atmospheric Administration, Boulder, Colorado, USA.

³Now at Aerodyne Research, Inc., Billerica, Massachusetts, USA.

⁴Now at Nutritional Science and Toxicology Department, University of California, Berkeley, California, USA.

⁵Laboratory of Atmospheric Chemistry, Paul Scherrer Institut, Villigen, Switzerland.

⁶Now at Science and Technology Corporation, Boulder, Colorado, USA.

⁷Department of Chemistry and Biochemistry and Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland.

⁸School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, Georgia, USA.




Figure 1. (a) Correlation plot of OA vs. CO in relatively fresh (toluene/benzene = 2–2.4) and aged (toluene/benzene < 1) plumes. (b, c) Average fossil and non-fossil aerosol TC mass from radiocarbon and Sunset OC/EC measurements as well as diurnal profiles of benzene and acetaldehyde measurements at Pasadena on two weekdays during CalNex.

[4] In this study, we first establish that in the LA Basin SOA makes a major contribution to OA mass and that its formation is dominated by fossil sources of carbon. We then use day of the

week differences in diesel and gasoline emissions and quantify the relative importance of gasoline and diesel emissions to secondary formation of OA by comparing airborne measurements of OA in the LA Basin between weekends and weekdays. We analyze data collected on 3 weekday (4 May, 14 May, and 19 May 2010) and 3 weekend (8 May (Saturday), 16 May (Sunday), and 20 June (Sunday) 2010) aircraft flights conducted in the LA Basin during daytime, as part of the CalNex (“Research at the Nexus of Air Quality and Climate Change”) field project.

2. Measurements

[5] During May and June 2010, the NOAA WP-3D aircraft conducted research flights in California as part of the CalNex field project. Here we analyze only those measurements which were collected in the LA Basin at altitudes lower than 1 km above sea level, between noon and 5:30 pm PDT (Figure S1 in Text S1). Measurements of submicron (50–650 nm) aerosol non-refractory composition aboard the aircraft were made using a compact time-of-flight aerosol mass spectrometer (C-ToF-AMS, referred to as AMS, Aerodyne Inc., Billerica, MA) [Bahreini *et al.*, 2009; Drewnick *et al.*, 2005]. Black carbon (BC) measurements in the size range of 90–700 nm were made by a single-particle soot photometer (SP2, Droplet Measurement Technologies, Inc., Boulder, CO) [Schwarz *et al.*, 2006]. Comparison of total mass estimated from the sum of AMS species and BC against total mass estimated from size distributions measured aboard the aircraft [Brock *et al.*, 2011] produces an average slope of 0.97 ($r^2 = 0.88$ and 0.97 for weekend and weekday, respectively) (Figure S2 in Text S1). Furthermore, the average weekday slope differs from the average weekend slope by less than 7%, indicating no significant instrumental bias between the weekend and weekday measurements.

[6] Gas phase measurements of volatile organic compounds (VOCs), including benzene, toluene, acetonitrile, formaldehyde, acetaldehyde, and isoprene aboard the aircraft were made online by a proton-transfer-reaction mass spectrometer (PTR-MS) [de Gouw and Warneke, 2007; Warneke *et al.*, 2011]. CO mixing ratios were measured by vacuum UV resonance fluorescence [Holloway *et al.*, 2000]. Mixing ratios of nitrogen dioxide and ozone were measured by a cavity ring-down spectrometer (CRDS) [Wagner *et al.*, 2011]. Photolysis rates of various species, including ozone and formaldehyde, were calculated using measurements from an actinic flux spectroradiometer aboard the aircraft [Stark *et al.*, 2007]. All aircraft measurements were averaged to the 10 s sampling time of the AMS.

[7] In addition to the airborne measurements during CalNex, ground-based measurements of gas and aerosol phase species were made in Pasadena, 17 km NE of downtown LA, during 15 May–15 June, 2010. At this site, total particulate carbon (TC) measurements were made by an EC/OC analyzer (Sunset Laboratory Inc., Tigard, OR) using the EUSAAR2 analytical method [Cavalli *et al.*, 2010]. Radiocarbon measurements were also performed offline on aerosol filter samples, collected every 3–4 h by a Digital DHA-80 HiVol sampler (DIGITEL Elektronik AG, Hegnau, Switzerland) [Szidat *et al.*, 2004]. Furthermore, VOC measurements of benzene and acetaldehyde were made online by GC-MS analysis [Goldan *et al.*, 2004].

3. Observations and Discussion

3.1. Anthropogenic Contribution to OA

[8] Airborne measurements of OA in the LA Basin show a significant (factor of 3) increase in the enhancement ratio of




Figure 2. Weekday (black circles) and weekend (orange circles) correlation plots of (a, b) OA vs. CO, (c) NO_x vs. CO, (d) BC vs. CO, (e) benzene vs. CO, and (f) toluene vs. benzene. Data were limited to altitudes lower than 1000 m over the LA Basin, and obtained between noon and 5:30 pm PDT. The dashed lines in Figures 2a and 2b indicate estimated urban $\Delta POA/\Delta CO$ [de Gouw *et al.*, 2008].

OA with respect to CO ($\Delta OA/\Delta CO$) with an increase in photochemical processing (i.e., decrease in the ratio of toluene/benzene) (Figure 1a). Consistent with previous studies [Docherty *et al.*, 2008; Turpin and Huntzicker, 1991; Williams *et al.*, 2010], this indicates a dominant contribution of SOA to the observed OA in the Basin. Furthermore, diurnal profiles of TC measurements in Pasadena reveal that there is a significant increase in TC mass coincident with the increase in secondary pollutants like acetaldehyde when the processed emissions from LA arrive at Pasadena (Figures 1b and 1c). Considering that BC was <15% of the TC mass (Figure 2d), TC closely represents OC during CalNex. In addition, the afternoon increase in TC mass is predominantly due to the increase in the fossil fraction, suggesting that photooxidation of anthropogenic organic compounds is the major contributor to SOA within the city plume [Zhang *et al.*, 2011].

[9] Lack of a significant contribution from biogenic VOCs to OA mass in the LA Basin is also apparent in the long-term measurements (<http://www.arb.ca.gov/aqd/aqcd/aqcdcdld.htm>) of OC and CO, a marker for primary combustion emissions, since anthropogenic emissions have decreased over the last decades with technological improvements in mobile sources [Ban-Weiss *et al.*, 2008] while the biogenic emissions are not expected to show a significant long-term trend. These measurements indicate that CO and OC have decreased by 3–6% per year in the last 10 years whereas OC/CO ratios have not decreased significantly (Figure S3 in Text S1). In addition, comparison of airborne fine particle volume measured from the NOAA P3 research aircraft on the west side of the LA basin on 13 May 2002 (a Monday) with similar measurements in 2010 indicates that the ratio of $\Delta \text{volume}/\Delta CO$ has stayed the same within the measurement uncertainties despite a factor of two reduction in CO and VOCs (not shown) from 2002 to 2010 (Figure S4 in Text S1). Considering the

small contribution of biomass burning (confirmed by low acetonitrile mixing ratios of 90 ± 22 pptv) to the 2010 data set, the above observations indicate that OA in the LA Basin is controlled by secondary processing of anthropogenic emissions.

3.2. Differences in Weekday-Weekend Enhancement Ratios

[10] Because of the different emissions from gasoline and diesel sources on weekends compared to weekdays, we use the airborne weekday-weekend data to distinguish the contribution of gasoline and diesel emissions to OA formation. On both weekdays and weekends, enhancements of OA with respect to CO (ratio of $\Delta\text{OA}/\Delta\text{CO}$) were much higher than the literature values of the enhancement ratios of primary OA ($\Delta\text{POA}/\Delta\text{CO}$) [de Gouw *et al.*, 2008], demonstrating the importance of secondary production of OA on both weekdays and weekends (Figures 2a and 2b). Consistent with a reduction in diesel emissions on weekends [Pollack *et al.*, 2012], $\Delta\text{NO}_x/\Delta\text{CO}$ is decreased by $63 \pm 5\%$ on weekends compared to weekdays (Figure 2c, uncertainties represent standard deviations of the linear least-square regression fits). Similarly, the $\Delta\text{BC}/\Delta\text{CO}$ in the LA Basin is decreased by $47 \pm 3\%$ on weekends compared to weekdays (Figure 2d). Consistent with other observations in the South Coast Air Basin, average weekend CO mixing ratios in Pasadena during CalNex are lower by only $4 \pm 11\%$ compared to weekday values [Pollack *et al.*, 2012]. Since nearly all of the CO emissions are from gasoline vehicles [Harley *et al.*, 2001], these observations are consistent with traffic data showing that gasoline emissions on weekends do not differ significantly from weekdays. Contrary to NO_x , BC does not undergo chemical loss in the atmosphere. Thus, using BC enhancement ratios (Figure 2d), emission factors of BC from gasoline and diesel vehicles [Ban-Weiss *et al.*, 2008], and fuel-use data, we estimate that there is a $54 \pm 5\%$ average reduction in diesel emissions on weekends compared to weekdays (see auxiliary material). Furthermore, we estimate that diesel vehicles contribute $87 \pm 1\%$ and $76 \pm 3\%$ of BC during weekdays and weekends, respectively. Note that these estimates are based on BC emission factors from studies in 2006. With the new particle filter requirements on diesel trucks traveling through the ports in California, BC emission factor from the modified diesel trucks may be lower by 50% [Dallmann *et al.*, 2011]. If these new emission factors represent the basin-wide diesel emissions, diesel contribution to BC during the week will be lower ($77 \pm 2\%$). Therefore, the analysis presented here depicts the upper estimate for diesel contributions.

3.3. Effects of Photochemical Processing on SOA Formation

[11] The 57% higher $\Delta\text{OA}/\Delta\text{CO}$ ratio observed on the weekends (Figures 2a and 2b) indicates that OA production per CO emitted is higher on the weekends despite the smaller contribution from diesel emissions. A question that arises is what fraction of the higher SOA on weekends is due to changes in photochemistry resulting from the lower NO_x emissions as opposed to changes in emissions on the weekends. The enhancement ratios of benzene with respect to CO are similar regardless of the day of the week (Figure 2e). Consistent with previous studies, this indicates that gasoline vehicles are the dominant source of CO and light, single-ring-aromatic VOCs including benzene and toluene [Marr and Harley, 2002b; Warneke *et al.*, 2007]. Because of the higher reactivity of toluene with OH radicals compared to benzene, the lower ratio of

toluene/benzene observed on the weekends (Figure 2f) confirms that faster photochemical processing of VOCs occurs during the weekends [Pollack *et al.*, 2012]. The actual rate of photochemical processing is not important for this analysis; however, the relative speed of photochemical processing due to OH during the weekends compared to weekdays is estimated by two independent methods (see auxiliary material). Both methods indicate that the rate of photochemical processing is higher on weekends, by factors of 1.9 ± 1.2 and 2.8 ± 1.0 . High VOC/ NO_x ratios can also increase SOA production yields [Ng *et al.*, 2007]; however, assuming 25 pptv of HO_2 , >95% of the observed weekend NO mixing ratios are still above the threshold (~ 20 pptv NO) below which SOA yields are expected to increase significantly [Henze *et al.*, 2008].

[12] To separate the role of faster photochemistry from emission differences on the weekends compared to weekdays, we use observed ratios of toluene/benzene to segregate OA and CO data by different degrees of photochemical processing, with the highest ratios indicative of the least processing (Figure 3). Lack of a significant difference in the slopes of the fitted lines to OA vs. CO in each panel indicates that similar enhancements in OA are observed on weekends and weekdays, for similar degrees of photochemical processing, despite the lower diesel emissions on weekends. Further, $\Delta\text{OA}/\Delta\text{CO}$ ratios for the least processed plumes (toluene/benzene = 2–2.4) are most similar to literature $\Delta\text{POA}/\Delta\text{CO}$ ratios while $\Delta\text{OA}/\Delta\text{CO}$ ratios increase by a factor of 1.5 for the plumes with toluene/benzene = 1–1.6.

3.4. Diesel Exhaust Contribution to SOA Formation

[13] The similar OA enhancements on weekends, when diesel emissions are low, allow us to calculate an upper limit to the contribution of diesel emissions to SOA formation. We assume that the POA mass does not change significantly with photochemical processing and subtract the estimated urban ratio of $\Delta\text{POA}/\Delta\text{CO}$ ($0.009 \pm 0.002 \mu\text{g m}^{-3} \text{ ppbv}^{-1}$ [de Gouw *et al.*, 2008]) from the observed $\Delta\text{OA}/\Delta\text{CO}$ ratios in Figure 3. Using the subtracted values, we calculate the ratio of weekday to weekend $\Delta\text{SOA}/\Delta\text{CO}$ for each photochemical processing category shown in Figure 3. The average ratio of weekday to weekend $\Delta\text{SOA}/\Delta\text{CO}$, weighted by the propagated uncertainties, is 0.90 ± 0.43 , which indicates that on average the contribution from diesel emissions to SOA formation is zero within the uncertainties. Using the estimate of $\Delta\text{POA}/\Delta\text{CO}$ [de Gouw *et al.*, 2008], POA emission factors [Ban-Weiss *et al.*, 2008], and fuel use data, we estimate that diesel engine emissions contribute $70 \pm 2\%$ of the POA in the LA Basin. Because diesel emissions contribute to POA, but not detectably to SOA, as photochemical processing and SOA formation proceeds, the contribution of diesel emissions to total OA decreases [Chirico *et al.*, 2010].

[14] We use the standard deviation of the weekday to weekend ratio of $\Delta\text{SOA}/\Delta\text{CO}$ to estimate an upper limit of 47% for the contribution of diesel engine emissions to total SOA on weekdays in the LA Basin and a lower limit of zero (see auxiliary material). Note that because of the lower diesel emissions on the weekends, weekend $\Delta\text{POA}/\Delta\text{CO}$ is most likely lower than the value assumed here. An alternative calculation that uses a lower anthropogenic $\Delta\text{POA}/\Delta\text{CO}$ based on the data from the CalNex ground site produces a maximum weekday contribution from diesel engine emissions to SOA of 20% (see auxiliary material). The estimates presented here represent conservative upper limits; on average diesel contribution to SOA is zero within the uncertainties. We cannot detect a diesel engine signature in the SOA measurements in




Figure 3. Correlation plots of OA vs. CO for different ratios of toluene to benzene (Tol./Benz.).

the LA Basin, while in contrast weekday enhancements in BC from diesel engine emissions are clearly evident.

4. Implications

[15] Our study suggests that, for more accurate modeling of SOA formation in urban areas, future research should be directed at identifying specific species in the exhaust of gasoline engines that are responsible for SOA formation. Studies targeted at understanding SOA formation from mobile sources with different after-treatment technologies would also be valuable. Our analysis suggests that the ratio of SOA to POA from gasoline emissions may reach a factor of ~ 3 at OH exposure of 5×10^7 molecule cm^{-3} h. A global estimate of primary OC (POC) from on-road gasoline emissions is 904 Gg/yr [Bond et al., 2004]. Assuming that production of SOA relative to POA from gasoline exhaust follows the same trend as in LA, and

using a ratio of 1.2 for POA/POC [Aiken et al., 2008], we estimate that within a day of processing, SOA from gasoline exhaust emissions may reach 4 Tg/yr, which is $\sim 16\%$ of recent global estimates of biogenic SOA [Andreae and Rosenfeld, 2008; Monks et al., 2009]. Our observations suggest that a decrease in the emission of organic species from gasoline engines may significantly reduce SOA concentrations on local and global scales.

[16] **Acknowledgments.** NOAA's Climate Change and Air Quality Programs supported this work. California Air Resources Board supported infrastructure at the Pasadena ground-site. NSF supported travel/Shipping costs of the PSI group. Access to the instrument MICADAS for ^{14}C analysis was provided by L. Wacker (ETH Zurich). We thank members of J. Surratt's and R. Weber's groups for support during the off-line sampling in Pasadena and I.B. Pollack for sharing results of her paper before publication. Discussions and feedback from J. L. Jimenez, A.L. Robinson, and I. B. Pollack are appreciated.

[17] The Editor wishes to thank two anonymous reviewers for their assistance evaluating this paper.

References

- Aiken, A. C., et al. (2008), O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with high-resolution time-of-flight aerosol mass spectrometry, *Environ. Sci. Technol.*, **42**, 4478–4485, doi:10.1021/es703009q.
- Andreae, M. O., and D. Rosenfeld (2008), Aerosol–cloud precipitation interactions. Part 1. The nature and sources of cloud active aerosols, *Earth Sci. Rev.*, **89**, 13–41, doi:10.1016/j.earscirev.2008.03.001.
- Bahreini, R., et al. (2009), Organic aerosol formation in urban and industrial plumes near Houston and Dallas, Texas, *J. Geophys. Res.*, **114**, D00F16, doi:10.1029/2008JD011493.
- Ban-Weiss, G. A., et al. (2008), Long-term changes in emissions of nitrogen oxides and particulate matter from on-road gasoline and diesel vehicles, *Atmos. Environ.*, **42**, 220–232, doi:10.1016/j.atmosenv.2007.09.049.
- Bond, T. C., D. G. Streets, K. F. Yarber, S. M. Nelson, J.-H. Woo, and Z. Klimont (2004), A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.*, **109**, D14203, doi:10.1029/2003JD003697.
- Brock, C. A., D. M. Murphy, R. Bahreini, and A. M. Middlebrook (2011), Formation and growth of organic aerosols downwind of the Deepwater Horizon oil spill, *Geophys. Res. Lett.*, **38**, L17805, doi:10.1029/2011GL048541.
- Cavalli, F., et al. (2010), Toward a standardized thermal-optical protocol for measuring atmospheric organic and elemental carbon: The EUSAAR protocol, *Atmos. Meas. Tech.*, **3**, 79–89, doi:10.5194/amt-3-79-2010.
- Chirico, R., et al. (2010), Impact of aftertreatment devices on primary emissions and secondary organic aerosol formation potential from in-use diesel vehicles: Results from smog chamber experiments, *Atmos. Chem. Phys.*, **10**, 11,545–11,563, doi:10.5194/acp-10-11545-2010.
- Dallmann, T. R., et al. (2011), Effects of diesel particle filter retrofits and accelerated fleet turnover on dryage truck emissions at the port of Oakland, *Environ. Sci. Technol.*, **45**, 10,773–10,779, doi:10.1021/es202609q.
- de Gouw, J. A., and C. Warneke (2007), Measurements of volatile organic compounds in the Earth's atmosphere using proton-transfer-reaction mass spectrometry, *Mass Spectrom. Rev.*, **26**, 223–257, doi:10.1002/mas.20119.
- de Gouw, J. A., et al. (2008), Sources of particulate matter in the northeastern United States in summer: 1. Direct emissions and secondary formation of organic matter in urban plumes, *J. Geophys. Res.*, **113**, D08301, doi:10.1029/2007JD009243.
- Docherty, K. S., et al. (2008), Apportionment of primary and secondary organic aerosols in Southern California during the 2005 Study of Organic Aerosols in Riverside (SOAR-1), *Environ. Sci. Technol.*, **42**, 7655–7662, doi:10.1021/es8008166.
- Drewnick, F., et al. (2005), A new time-of-flight aerosol mass spectrometer (TOF-AMS)—Instrument description and first field deployment, *Aerosol Sci. Technol.*, **39**, 637–658, doi:10.1080/02786820500182040.
- Fuzzi, S., et al. (2006), Critical assessment of the current state of scientific knowledge, terminology, and research needs concerning the role of organic aerosols in the atmosphere, climate, and global change, *Atmos. Chem. Phys.*, **6**, 2017–2038, doi:10.5194/acp-6-2017-2006.
- Goldan, P. D., W. C. Kuster, E. Williams, P. C. Murphy, F. C. Fehsenfeld, and J. Meagher (2004), Nonmethane hydrocarbon and oxy hydrocarbon measurements during the 2002 New England Air Quality Study, *J. Geophys. Res.*, **109**, D21309, doi:10.1029/2003JD004455.
- Griffin, R. J., et al. (1999), Organic aerosol formation from the oxidation of biogenic hydrocarbons, *J. Geophys. Res.*, **104**, 3555–3567, doi:10.1029/1998JD100049.

- Harley, R. A., S. A. McKeen, J. Pearson, M. O. Rodgers, and W. A. Lonneman (2001), Analysis of motor vehicle emissions during the Nashville/Middle Tennessee Ozone Study, *J. Geophys. Res.*, **106**, 3559–3567, doi:10.1029/2000JD900677.
- Harley, R. A., et al. (2005), Changes in motor vehicle emissions on diurnal to decadal time scales and effects on atmospheric composition, *Environ. Sci. Technol.*, **39**, 5356–5362, doi:10.1021/es048172+.
- Henze, D. K., et al. (2008), Global modeling of secondary organic aerosol formation from aromatic hydrocarbons: High- vs. low-yield pathways, *Atmos. Chem. Phys.*, **8**, 2405–2420, doi:10.5194/acp-8-2405-2008.
- Holloway, J. S., R. O. Jakoubek, D. D. Parrish, C. Gerbig, A. Volz-Thomas, S. Schmitgen, A. Fried, B. Wert, B. Henry, and J. R. Drummond (2000), Airborne intercomparison of vacuum ultraviolet fluorescence and tunable diode laser absorption measurements of tropospheric carbon monoxide, *J. Geophys. Res.*, **105**, 24,251–24,261, doi:10.1029/2000JD900237.
- Kleeman, M. J., et al. (2008), Lubricating oil and fuel contributions to particulate matter emissions from light-duty gasoline and heavy-duty diesel vehicles, *Environ. Sci. Technol.*, **42**, 235–242, doi:10.1021/es071054c.
- Marr, L. C., and R. A. Harley (2002a), Modeling the effect of weekday-weekend differences in motor vehicle emissions on photochemical air pollution in central California, *Environ. Sci. Technol.*, **36**, 4099–4106, doi:10.1021/es020629x.
- Marr, L. C., and R. A. Harley (2002b), Spectral analysis of weekday-weekend differences in ambient ozone, nitrogen oxide, and non-methane hydrocarbon time series in California, *Atmos. Environ.*, **36**, 2327–2335, doi:10.1016/S1352-2310(02)00188-7.
- Miracolo, M. A., et al. (2010), Photo-oxidation of low-volatility organics found in motor vehicle emissions: Production and chemical evolution of organic aerosol mass, *Environ. Sci. Technol.*, **44**, 1638–1643, doi:10.1021/es902635c.
- Monks, P. S., et al. (2009), Atmospheric composition change—Global and regional air quality, *Atmos. Environ.*, **43**, 5268–5350, doi:10.1016/j.atmosenv.2009.08.021.
- Murphy, D. M., et al. (2008), Weekly patterns of aerosol in the United States, *Atmos. Chem. Phys.*, **8**, 2729–2739, doi:10.5194/acp-8-2729-2008.
- Murphy, J. G., et al. (2007), The weekend effect within and downwind of Sacramento—Part 1: Observations of ozone, nitrogen oxides, and VOC reactivity, *Atmos. Chem. Phys.*, **7**, 5327–5339, doi:10.5194/acp-7-5327-2007.
- Ng, N. L., et al. (2007), Secondary organic aerosol formation from m-xylene, toluene, and benzene, *Atmos. Chem. Phys.*, **7**, 3909–3922, doi:10.5194/acp-7-3909-2007.
- Odum, J. R., et al. (1997), The atmospheric aerosol-forming potential of whole gasoline vapor, *Science*, **276**, 96–99, doi:10.1126/science.276.5309.96.
- Pollack, I. B., et al. (2012), Airborne and ground-based observations of a weekend effect in ozone, precursors, and oxidation products in the California South Coast Air Basin, *J. Geophys. Res.*, **117**, D00V05, doi:10.1029/2011JD016772.
- Presto, A. A., et al. (2009), Intermediate-volatility organic compounds: A potential source of ambient oxidized organic aerosol, *Environ. Sci. Technol.*, **43**, 4744–4749, doi:10.1021/es803219q.
- Schwarz, J. P., et al. (2006), Single-particle measurements of midlatitude black carbon and light-scattering aerosols from the boundary layer to the lower stratosphere, *J. Geophys. Res.*, **111**, D16207, doi:10.1029/2006JD007076.
- Stark, H., B. M. Lerner, R. Schmitt, R. Jakoubek, E. J. Williams, T. B. Ryerson, D. T. Sueper, D. D. Parrish, and F. C. Fehsenfeld (2007), Atmospheric in situ measurement of nitrate radical (NO₃) and other photolysis rates using spectroradiometry and filter radiometry, *J. Geophys. Res.*, **112**, D10S04, doi:10.1029/2006JD007578.
- Szidat, S., et al. (2004), Source apportionment of aerosols by C-14 measurements in different carbonaceous particle fractions, *Radiocarbon*, **46**, 475–484.
- Turpin, B. J., and J. J. Huntzicker (1991), Secondary formation of organic aerosol in the Los Angeles Basin: A descriptive analysis of organic and elemental carbon concentrations, *Atmos. Environ.*, **25A**(2), 207–215.
- Wagner, N. L., et al. (2011), Diode laser-based cavity ring-down instrument for NO₃, N₂O₅, NO, NO₂, and O₃ from aircraft, *Atmos. Meas. Tech.*, **4**(6), 1227–1240, doi:10.5194/amt-4-1227-2011.
- Warneke, C., et al. (2007), Determination of urban volatile organic compound emission ratios and comparison with an emissions database, *J. Geophys. Res.*, **112**, D10S47, doi:10.1029/2006JD007930.
- Warneke, C., et al. (2011), Airborne formaldehyde measurements using PTR-MS: Calibration, humidity dependence, inter-comparison and initial results, *Atmos. Meas. Tech.*, **4**, 2345–2358, doi:10.5194/amt-4-2345-2011.
- Williams, B. J., et al. (2010), Major components of atmospheric organic aerosol in Southern California as determined by hourly measurements of source marker compounds, *Atmos. Chem. Phys.*, **10**, 11,577–11,603, doi:10.5194/acp-10-11577-2010.
- Zhang, Q., et al. (2007), Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophys. Res. Lett.*, **34**, L13801, doi:10.1029/2007GL029979.
- Zhang, X., Y.-H. Lin, J. D. Surratt, P. Zotter, A. S. H. Prevôt, and R. J. Weber (2011), Light-absorbing soluble organic aerosol in Los Angeles and Atlanta: A contrast in secondary organic aerosol, *Geophys. Res. Lett.*, **38**, L21810, doi:10.1029/2011GL049385.
- R. Bahreini, C. A. Brock, S. S. Brown, J. A. de Gouw, W. P. Dube, J. B. Gilman, J. S. Holloway, W. C. Kuster, A. M. Middlebrook, D. D. Parrish, A. E. Perring, J. P. Schwarz, M. Trainer, N. L. Wagner, and C. Warneke, Earth System Research Laboratory, National Oceanic and Atmospheric Administration, 325 Broadway, Boulder, CO 80305, USA. (roya.bahreini@noaa.gov)
- K. Hall, Nutritional Science and Toxicology Department, University of California, 119 Morgan Hall #3104, Berkeley, CA 94720, USA.
- A. S. H. Prevôt and P. Zotter, Laboratory of Atmospheric Chemistry, Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland.
- J. R. Spackman, Science and Technology Corporation, Boulder, Colorado, USA.
- H. Stark, Aerodyne Research, Inc., 45 Manning Rd., Billerica, MA 01821, USA.
- S. Szidat, Department of Chemistry and Biochemistry, University of Bern, Freiestrasse 3, CH-3012 Bern, Switzerland.
- R. J. Weber, School of Earth and Atmospheric Sciences, Georgia Institute of Technology, 311 Ferst Dr., Atlanta, GA 30332, USA.